

## Matricial Logarithmic Derivatives\*

Emeric Deutsch

*Department of Mathematics  
Polytechnic Institute of New York  
Brooklyn, New York 11201*

and

Max Mlynarski<sup>†</sup>

*Department of Mathematics  
Kingsborough Community College  
Brooklyn, New York 11235*

Submitted by Hans Schneider

---

### ABSTRACT

If  $\phi$  is a norm on  $\mathbf{C}^n$ , then the mapping  $A \rightarrow \lim_{h \downarrow 0} [|I + hA|_\phi - 1]/h$  from  $M_n(\mathbf{C})$  ( $= \mathbf{C}^{n \times n}$ ) into  $\mathbf{R}$  is called the logarithmic derivative induced by the vector norm  $\phi$ . In this paper we generalize this concept to a mapping  $\gamma$  from  $M_n(\mathbf{C})$  into  $M_k(\mathbf{R})$ , where  $k \leq n$ . Denoting by  $\alpha(B)$  the spectral abscissa of a square matrix  $B$  (the largest of the real parts of the eigenvalues), we show, in particular, that  $\alpha(A) \leq \alpha(\gamma(A))$ . As a byproduct we obtain simple sufficient conditions for the stability of a matrix.

---

### 1. INTRODUCTION

Let  $\mathbf{C}^n$  denote the vector space of column  $n$ -tuples of complex numbers, and let  $M_n(\mathbf{C})$  denote the algebra of complex  $n \times n$  matrices. If  $\phi: \mathbf{C}^n \rightarrow \mathbf{R}$  is a norm on  $\mathbf{C}^n$  and  $\text{lub}_\phi$  denotes the corresponding operator norm on  $M_n(\mathbf{C})$ , then the mapping

$$\gamma_\phi: M_n(\mathbf{C}) \rightarrow \mathbf{R},$$
$$\gamma_\phi(A) = \lim_{h \downarrow 0} \frac{\text{lub}_\phi(I_n + hA) - 1}{h} \quad [A \in M_n(\mathbf{C})] \quad (1.1)$$

---

\*Parts of this paper were presented at the Seminar on Advances in Matrix Theory (Positivity and Norms), held in München, December 10–13, 1974.

<sup>†</sup>This research is part of the dissertation submitted to the Faculty of the Polytechnic Institute of New York in partial fulfillment of the requirements for the degree of Doctor of Philosophy, awarded to Max Mlynarski in 1976.

( $I_n$  is the  $n \times n$  identity matrix) is called *the logarithmic derivative induced by the vector norm  $\phi$* . It is also called the logarithmic norm, although it is not a norm. The concept of logarithmic derivative has been introduced separately by Dahlquist [2, p. 10] and Lozinskii [15] as a tool to study the growth of solutions of ordinary differential equations (see also Coppel [1, p. 41]).

In the case of the Hölder norms  $l_2, l_\infty, l_1$ , we have for  $A = (a_{ij}) \in M_n(\mathbb{C})$ ,

$$\gamma_{l_2}(A) = \text{largest eigenvalue of } \frac{1}{2}(A + A^*), \quad (1.2)$$

$$\gamma_{l_\infty}(A) = \max_i \left[ \operatorname{Re} a_{ii} + \sum_{\substack{j=1 \\ j \neq i}}^n |a_{ij}| \right], \quad (1.3)$$

$$\gamma_{l_1}(A) = \max_j \left[ \operatorname{Re} a_{jj} + \sum_{\substack{i=1 \\ i \neq j}}^n |a_{ij}| \right]. \quad (1.4)$$

The basic properties of the mapping  $\gamma_\phi$  are:

- (i)  $\gamma_\phi(A + B) \leq \gamma_\phi(A) + \gamma_\phi(B) \quad \forall A, B \in M_n(\mathbb{C})$ ;
- (ii)  $\gamma_\phi(\zeta A) = \begin{cases} \zeta \gamma_\phi(A) & \forall \zeta \geq 0, \quad \forall A \in M_n(\mathbb{C}), \\ -\zeta \gamma_\phi(-A) & \forall \zeta \leq 0, \quad \forall A \in M_n(\mathbb{C}); \end{cases}$
- (iii)  $\gamma_\phi(A + \zeta I_n) = \gamma_\phi(A) + \operatorname{Re} \zeta \quad \forall \zeta \in \mathbb{C}, \quad \forall A \in M_n(\mathbb{C})$ ;
- (iv)  $|\gamma_\phi(A)| \leq \operatorname{lub}_\phi A \quad \forall A \in M_n(\mathbb{C})$ ;
- (v)  $\operatorname{Re} \lambda \leq \gamma_\phi(A)$  for each eigenvalue  $\lambda$  of  $A$ ,  $\quad \forall A \in M_n(\mathbb{C})$ .

The purpose of this paper is to introduce and investigate a generalization of the concept of logarithmic derivative induced by a vector norm.

## 2. NOTATIONS, TERMINOLOGY AND PRELIMINARIES

(i)

We denote by  $\mathbf{C}^n(\mathbf{R}^n)$  the vector space of all column  $n$ -tuples of complex (real) numbers. The vector space  $\mathbf{R}^n$  is partially ordered componentwise. We denote  $\mathbf{R}_+^n = \{x \in \mathbf{R}^n : x \geq 0\}$ . The vectors of the standard basis of  $\mathbf{C}^n$  will be denoted by  $e_1, e_2, \dots, e_n$ .

A finite collection of subspaces  $\pi = \{W_1, \dots, W_k\}$  is said to be a *direct-sum decomposition* of  $\mathbf{C}^n$  if  $\mathbf{C}^n = W_1 \oplus W_2 \oplus \dots \oplus W_k$ . In particular, if

$$W_i = \operatorname{span}\{e_{\eta_{i-1}+1}, e_{\eta_{i-1}+2}, \dots, e_{\eta_i}\}$$

( $j=1, \dots, k$ ), where  $r_0, r_1, \dots, r_k$  are nonnegative integers satisfying

$$0 = r_0 < r_1 < \dots < r_k = n, \quad (2.1)$$

then we say that  $\pi = \{W_1, \dots, W_k\}$  is a *partition* of  $\mathbf{C}^n$ . Clearly, a partition  $\pi$  of  $\mathbf{C}^n$  is completely determined by integers  $r_0, r_1, \dots, r_k$  satisfying (2.1). In this case we shall write  $\pi = \{r_0, r_1, \dots, r_k\}$ .

(ii)

We denote by  $M_n(\mathbf{C})$  [ $M_n(\mathbf{R})$ ] the algebra of all complex [real]  $n \times n$  matrices. The  $n \times n$  identity matrix will be denoted by  $I_n$ .

Let  $A \in M_n(\mathbf{C})$ . We shall denote by  $\Lambda(A)$  the *spectrum* of  $A$ , by  $r(A)$  the *spectral radius* of  $A$ , i.e.,

$$r(A) = \max\{|\lambda| : \lambda \in \Lambda(A)\},$$

and by  $\alpha(A)$  the *spectral abscissa* of  $A$ , i.e.,

$$\alpha(A) = \max\{\operatorname{Re} \lambda : \lambda \in \Lambda(A)\}.$$

It can be easily seen that  $r(e^A) = e^{\alpha(A)}$  and that

$$\alpha(A) = \lim_{h \downarrow 0} \frac{r(I_n + hA) - 1}{h} = \inf_{h > 0} \frac{r(I_n + hA) - 1}{h} \quad (2.2)$$

for all  $A \in M_n(\mathbf{C})$ .

A matrix  $A \in M_n(\mathbf{C})$  is said to be *stable* if  $\alpha(A) < 0$ , i.e., if all the eigenvalues of  $A$  lie in the open left half plane.

The algebra  $M_n(\mathbf{R})$  is partially ordered componentwise. If  $A = (a_{ij}) \in M_n(\mathbf{C})$ , we shall write  $|A| = (|a_{ij}|)$ .

A matrix  $A \in M_n(\mathbf{R})$  is said to be *nonnegative* if  $A \geq 0$ . The set of all nonnegative matrices in  $M_n(\mathbf{R})$  will be denoted by  $M_n(\mathbf{R}_+)$ . From the Perron-Frobenius theory of nonnegative matrices it is well known (see, for example, [24]) that

- (a)  $A \geq 0 \Rightarrow \alpha(A) = r(A) \in \Lambda(A)$ ,
- (b)  $0 \leq A \leq B \Rightarrow r(A) \leq r(B)$ .

A matrix  $A = (a_{ij}) \in M_n(\mathbf{R})$  is said to be *essentially nonnegative* if  $a_{ij} \geq 0$  for all  $i \neq j$ ,  $i, j = 1, 2, \dots, n$  (see [24, p. 260]). Since in this case  $A + \tau I_n$  is

nonnegative for sufficiently large  $\tau \in \mathbf{R}$ , it follows at once that

- (c)  $A$  essentially nonnegative  $\Rightarrow \alpha(A) \in \Lambda(A)$ ,
- (d)  $A, B$  essentially nonnegative,  $A \leq B \Rightarrow \alpha(A) \leq \alpha(B)$ .

A matrix  $A = (a_{ij}) \in M_n(\mathbf{R})$  is said to be an *M-matrix* [24, p. 85] if (1)  $a_{ij} \leq 0$  for  $i \neq j$ , (2)  $A$  is nonsingular, and (3)  $A^{-1} \geq 0$ .

(iii)

Let  $\phi$  be a vector norm on  $\mathbf{C}^n$ . The operator norm induced by  $\phi$  will be denoted by  $\text{lub}_\phi$  or by  $\|\cdot\|_\phi$ . It is also called *the matrix norm subordinate to  $\phi$*  or *the least upper bound norm associated with  $\phi$*  [12, p. 39].

If  $A, B \in M_n(\mathbf{C})$ , we denote

$$g_\phi(A, B) = \lim_{h \downarrow 0} \frac{\|A + hB\|_\phi - \|A\|_\phi}{h}. \quad (2.3)$$

It is known [14, p. 347] that this limit exists; it is called *the right Gateaux derivative of the norm  $\|\cdot\|_\phi$  ( $=\text{lub}_\phi$ ) at  $A$  with respect to  $B$* . The basic properties of  $g_\phi$  are

$$\begin{aligned} g_\phi(A, B_1 + B_2) &\leq g_\phi(A, B_1) + g_\phi(A, B_2), \\ g_\phi(A, \zeta B) &= \begin{cases} \zeta g_\phi(A, B) & \text{if } \zeta \geq 0, \\ -\zeta g_\phi(A, -B) & \text{if } \zeta \leq 0, \end{cases} \\ g_\phi(A, B + \zeta A) &= g_\phi(A, B) + \|A\|_\phi \text{Re } \zeta \quad \forall \zeta \in \mathbf{C}, \\ |g_\phi(A, B)| &\leq \|B\|_\phi \quad \forall A, B, B_1, B_2 \in M_n(\mathbf{C}). \end{aligned}$$

Obviously,  $\gamma_\phi(A) = g_\phi(I_n, A)$  for all  $A \in M_n(\mathbf{C})$ .

(iv)

A mapping  $p: \mathbf{C}^n \rightarrow \mathbf{R}_+^k$  satisfying

- (1)  $p(\zeta x) = |\zeta| p(x) \quad \forall x \in \mathbf{C}^n, \quad \forall \zeta \in \mathbf{C}$ ,
- (2)  $p(x + y) \leq p(x) + p(y) \quad \forall x, y \in \mathbf{C}^n$ ,
- (3)  $p(x) \neq 0 \quad \text{if } x \in \mathbf{C}^n, \quad x \neq 0$ ,

is called a *vectorial norm of order  $k$  on  $\mathbf{C}^n$*  [5, 13, 20, 21]. If  $\phi$  is a vector norm on  $\mathbf{C}^n$  and  $\pi = \{W_1, \dots, W_k\}$  is a direct-sum decomposition of  $\mathbf{C}^n$  with associated projections  $E_1, \dots, E_k$ , then the mapping  $p: \mathbf{C}^n \rightarrow \mathbf{R}_+^k$  defined by

$$p(x) = (\phi(E_1 x), \dots, \phi(E_k x)) \quad (x \in \mathbf{C}^n)$$

is a vectorial norm on  $\mathbf{C}^n$  [5]. We shall say that  $p$  is *induced by  $\phi$  and  $\pi$* , and

we shall write  $p = (\phi; \pi)$ . A vectorial norm obtained in this manner is said to be *regular*. (For equivalent definitions, see [5, 21]). A regular vectorial norm  $p = (\phi; \pi)$  is said to be *equilibrated* if  $\|E_j\|_\phi = 1$  for all  $j = 1, \dots, k$ , where the  $E_j$ 's are the projections associated with the direct-sum decomposition  $\pi$ .

If  $p$  is a vectorial norm on  $\mathbf{C}^n$  and  $G$  is a nonsingular matrix in  $M_n(\mathbf{C})$ , then the mapping  $x \rightarrow p(Gx)$  of  $\mathbf{C}^n$  into  $\mathbf{R}_+^k$  is also a vectorial norm on  $\mathbf{C}^n$ , called the *G-transform* of  $p$  and denoted by  $p_G$  [5].

(v)

A mapping  $\mu : M_n(\mathbf{C}) \rightarrow M_k(\mathbf{R}_+)$  satisfying

- (1)  $\mu(\zeta A) = |\zeta| \mu(A) \quad \forall A \in M_n(\mathbf{C}), \quad \forall \zeta \in \mathbf{C},$
- (2)  $\mu(A + B) \leq \mu(A) + \mu(B) \quad \forall A, B \in M_n(\mathbf{C}),$
- (3)  $\mu(AB) \leq \mu(A) \mu(B) \quad \forall A, B \in M_n(\mathbf{C}),$
- (4)  $\mu(A) \neq 0 \quad \text{if } A \in M_n(\mathbf{C}), \quad A \neq 0,$

is called a *matricial norm of order k* on  $M_n(\mathbf{C})$  [3].

Let  $p = (\phi; \pi)$  be a vectorial norm on  $\mathbf{C}^n$ , where  $\pi = \{W_1, \dots, W_k\}$ , and let  $E_1, \dots, E_k$  be the projections associated with  $\pi$ . Then, the mapping

$$\text{lub}_p : M_n(\mathbf{C}) \rightarrow M_n(\mathbf{R}_+),$$

defined by

$$\text{lub}_p A = (m_{ij}(A))_{i,j=1,\dots,k} \quad [A \in M_n(\mathbf{C})],$$

where

$$m_{ij}(A) = \sup_{\substack{x \in W_j \\ x \neq 0}} \frac{\phi(E_i A x)}{\phi(E_j x)},$$

is a matricial norm on  $M_n(\mathbf{C})$ , called the *matricial norm subordinate to the vectorial norm*  $p = (\phi; \pi)$  [4, 17, 20, 21]. This matrix is the least element of the set

$$\{B \in M_k(\mathbf{R}_+) : p(Ax) \leq Bp(x) \quad \forall x \in \mathbf{C}^n\}.$$

It is known [4] that in the case when  $p = (\phi; \pi)$  is equilibrated, then

$$\text{lub}_p A = (\|E_i A E_j\|_\phi)_{i,j=1,\dots,k}. \quad (2.4)$$

If  $A \in M_n(\mathbf{C})$ , and if  $p = (\phi; \pi)$  is a vectorial norm of order  $k$  on  $\mathbf{C}^n$ , then [3, 17, 21]

$$r(A) \leq r(\text{lub}_p A). \quad (2.5)$$

### 3. MATRICIAL LOGARITHMIC DERIVATIVES

DEFINITION 3.1. Let  $p$  be an equilibrated vectorial norm of order  $k$  on  $\mathbf{C}^n$ . The mapping

$$\gamma_p : M_n(\mathbf{C}) \rightarrow M_k(\mathbf{R}),$$

$$\gamma_p(A) = \lim_{h \downarrow 0} \frac{\text{lub}_p(I_n + hA) - I_k}{h} \quad [A \in M_n(\mathbf{C})]$$

is called *the matricial logarithmic derivative induced by  $p$* .

REMARK 3.1. If  $p = (\phi; \pi)$ , then

$$\frac{1}{h} [\text{lub}_p(I_n + hA) - I_k] = (\mu_{ij})_{i,j=1,\dots,k}, \quad (3.1)$$

where  $\mu_{ii} = [\|E_i + hE_iAE_i\|_\phi - 1]/h$  and  $\mu_{ij} = \|E_iAE_j\|_\phi$  ( $i \neq j$ ),  $E_1, \dots, E_k$  being the projections associated with the direct-sum decomposition  $\pi$ . Thus,  $\lim_{h \downarrow 0} \mu_{ii} = g_\phi(E_i, E_iAE_i)$ , and so the limit in the definition of  $\gamma_p(A)$  exists.

PROPOSITION 3.1. Let  $p = (\phi; \pi)$  be an equilibrated vectorial norm of order  $k$  on  $\mathbf{C}^n$ , and let  $E_1, \dots, E_k$  be the projections associated with  $\pi$ . Then,

(i)

$$\gamma_p(A) = \begin{pmatrix} g_\phi(E_1, E_1AE_1)_\phi & \|E_1AE_2\|_\phi & \cdots & \|E_1AE_k\|_\phi \\ \|E_2AE_1\|_\phi & g_\phi(E_2, E_2AE_2) & \cdots & \|E_2AE_k\|_\phi \\ \vdots & \vdots & \ddots & \vdots \\ \|E_kAE_1\|_\phi & \|E_kAE_2\|_\phi & \cdots & g_\phi(E_k, E_kAE_k) \end{pmatrix}, \quad (3.2)$$

(ii)  $\gamma_p(A)$  is an essentially nonnegative  $k \times k$  matrix.

*Proof.* The expression (3.2) follows at once from Remark 3.1, while (ii) follows from (3.2). ■

REMARK 3.2. Recently, matricial logarithmic derivatives have been introduced by Hewer [11], but only under the following restrictive assumption: it is assumed that the projections  $E_1, \dots, E_k$  associated with the direct-sum decomposition  $\pi$  commute with the matrix  $A$  for which the matricial logarithmic derivative is defined. It follows at once from (3.2) that in this case  $\gamma_p(A)$  is a diagonal matrix.

In [16], Lozinskii has investigated some matricial logarithmic derivatives for the special case when  $\phi$  is a Hölder norm and  $\pi$  is a partition of  $\mathbb{C}^n$ .

PROPOSITION 3.2. *If  $p$  is an equilibrated vectorial norm of order  $k$  on  $\mathbb{C}^n$ , then*

$$\gamma_p(A) = \inf_{h>0} \frac{\text{lub}_p(I_n + hA) - I_k}{h} \quad [A \in M_n(\mathbb{C})].$$

*Proof.* This follows at once from the fact that each entry in (3.1) is a nondecreasing function of  $h$  (see [14, pp. 347–348]). ■

PROPOSITION 3.3. *Let  $p$  be an equilibrated vectorial norm of order  $k$  on  $\mathbb{C}^n$ . Then,*

- (i)  $\gamma_p(A + B) \leq \gamma_p(A) + \gamma_p(B) \quad \forall A, B \in M_n(\mathbb{C});$
- (ii)  $\gamma_p(\xi A) = \begin{cases} \xi \gamma_p(A) & \forall \xi \geq 0, \quad \forall A \in M_n(\mathbb{C}), \\ -\xi \gamma_p(-A) & \forall \xi \leq 0, \quad \forall A \in M_n(\mathbb{C}); \end{cases}$
- (iii)  $\gamma_p(A + \xi I_n) = \gamma_p(A) + I_k \text{Re } \xi \quad \forall \xi \in \mathbb{C}, \quad \forall A \in M_n(\mathbb{C});$
- (iv)  $|\gamma_p(A)| \leq \text{lub}_p A \quad \forall A \in M_n(\mathbb{C}).$

*Proof.* All these properties follow from the corresponding properties of the right Gateaux derivative. To prove (iv), one has to take into account (2.4). ■

PROPOSITION 3.4. *Let  $p$  be an equilibrated vectorial norm of order  $k$  on  $\mathbb{C}^n$ . Then*

$$\text{lub}_p(e^A) \leq e^{\gamma_p(A)} \quad \forall A \in M_n(\mathbb{C}).$$

*Proof.* For each positive integer  $m$ , let

$$F_m = m \left[ \text{lub}_p \left( I_n + \frac{1}{m} A \right) - I_k \right] - \gamma_p(A).$$

Then  $F_m \rightarrow 0$  as  $m \rightarrow \infty$ . Now

$$\text{lub}_p \left[ \left( I_n + \frac{1}{m} A \right)^m \right] \leq \left[ \text{lub}_p \left( I_n + \frac{1}{m} A \right) \right]^m = \left( I_k + \frac{\gamma_p(A) + F_m}{m} \right)^m$$

and letting  $m \rightarrow \infty$ , we obtain the required inequality.  $\blacksquare$

**COROLLARY 3.1.** *Let  $p$  be an equilibrated vectorial norm of order  $k$  on  $\mathbb{C}^n$ , and let  $A \in M_n(\mathbb{C})$ . Then*

$$\text{lub}_p(e^{tA}) \leq e^{t\gamma_p(A)} \quad \forall t \geq 0.$$

*Proof.* This is an immediate consequence of Proposition 3.4 and Proposition 3.3(ii).  $\blacksquare$

**REMARK 3.3.** The inequality (iv) of Proposition 3.3 shows that the result of Corollary 3.1 is better than the easily proved inequality

$$\text{lub}_p(e^{tA}) \leq e^{t\text{lub}_p A} \quad \forall t \geq 0.$$

Moreover, the inequality of Corollary 3.1 cannot be improved, as is seen from the following proposition.

**PROPOSITION 3.5.** *Let  $p$  be an equilibrated vectorial norm of order  $k$  on  $\mathbb{C}^n$ , and let  $A \in M_n(\mathbb{C})$ . Then  $\gamma_p(A)$  is the least element of the set*

$$\{ S \in M_k(\mathbb{R}) : \text{lub}_p(e^{tA}) \leq e^{tS} \quad \forall t \geq 0 \}.$$

*Proof.* Denoting by  $\mathcal{K}$  the set given in the proposition, it follows at once from Proposition 3.4 that  $\gamma_p(A) \in \mathcal{K}$ . Now, assuming that  $S \in \mathcal{K}$ , we have to show that  $\gamma_p(A) \leq S$ . From  $\text{lub}_p(e^{tA}) \leq e^{tS}$  ( $t \geq 0$ ), we obtain

$$\text{lub}_p \left( I_n + \frac{t}{1!} A + \frac{t^2}{2!} A^2 + \cdots \right) \leq I_k + \frac{t}{1!} S + \frac{t^2}{2!} S^2 + \cdots \quad (t \geq 0),$$



whence

$$\text{lub}_p(I_n + tA) - I_k \leq tS + o(t) \quad (t \geq 0),$$

and so

$$\frac{1}{t} [\text{lub}_p(I_n + tA) - I_k] \leq S + \frac{1}{t} o(t) \quad (t \geq 0).$$

Letting  $t \downarrow 0$ , we obtain  $\gamma_p(A) \leq S$ . ■

#### 4. MATRICIAL LOGARITHMIC DERIVATIVES AND SPECTRAL ABSCISSAE

In this section we shall discuss various properties involving matricial logarithmic derivatives and spectral abscissae.

**PROPOSITION 4.1** *Let  $p$  be an equilibrated vectorial norm of order  $k$  on  $\mathbb{C}^n$ . Then*

$$\alpha(\gamma_p(A)) = \inf_{h>0} \frac{r(\text{lub}_p(I_n + hA)) - 1}{h} \quad [A \in M_n(\mathbb{C})].$$

*Proof.* Denote

$$B_h = \frac{1}{h} [\text{lub}_p(I_n + hA) - I_k] \quad (h > 0).$$

Since  $B_h$  is essentially nonnegative and the off-diagonal entries of  $B_h$  do not depend on  $h$  [see (3.1)], one can easily show that  $\inf_{h>0} \alpha(B_h) = \alpha(\inf_{h>0} B_h)$ . Then, making use of Proposition 3.2, we have

$$\begin{aligned} \alpha(\gamma_p(A)) &= \alpha\left(\inf_{h>0} B_h\right) = \inf_{h>0} \alpha(B_h) \\ &= \inf_{h>0} \frac{\alpha(\text{lub}_p(I_n + hA)) - 1}{h} = \inf_{h>0} \frac{r(\text{lub}_p(I_n + hA)) - 1}{h}. \end{aligned} \quad \blacksquare$$

PROPOSITION 4.2. *Let  $p$  be an equilibrated vectorial norm on  $\mathbf{C}^n$ . Then*

$$\alpha(A) \leq \alpha(\gamma_p(A)) \quad [A \in M_n(\mathbf{C})].$$

*Proof.* Making use of Proposition 3.4, we have

$$e^{\alpha(A)} = r(e^A) \leq r(\text{lub}_p e^A) \leq r(e^{\gamma_p(A)}) = e^{\alpha(\gamma_p(A))},$$

whence  $\alpha(A) \leq \alpha(\gamma_p(A))$ . ■

REMARK 4.1. An alternate proof of Proposition 4.2 goes as follows: we have, for all  $h \geq 0$  [see (2.5)],

$$1 + h\alpha(A) = \alpha(I_n + hA) \leq r(I_n + hA) \leq r(\text{lub}_p(I_n + hA)),$$

whence

$$\alpha(A) \leq \frac{1}{h} [r(\text{lub}_p(I_n + hA)) - 1] \quad (h > 0).$$

Letting  $h \downarrow 0$  and making use of Proposition 4.1, we obtain  $\alpha(A) \leq \alpha(\gamma_p(A))$ .

REMARK 4.2. Proposition 4.2 has been proved by Deutsch [6] by a more cumbersome method and only for the special case when  $p = (\phi; \pi)$ ,  $\pi$  being a partition of  $\mathbf{C}^n$ .

EXAMPLE 4.1. Let

$$A = \begin{pmatrix} 8 & 1 & 2 & -5 & -1 \\ 1 & 1 & -2 & 0 & 0 \\ 1 & 1 & 0 & -2 & 0 \\ 0 & 1 & 0 & 0 & -4 \end{pmatrix},$$

and consider the partition  $\pi = \{0, 1, 4\}$  of  $\mathbf{C}^4$ , which induces the indicated partitioning of  $A$ . Denoting  $p = (I_\infty, \pi)$ , we have

$$\gamma_p(A) = \begin{pmatrix} 8 & 8 \\ 1 & -2 \end{pmatrix},$$

and so  $\alpha(\gamma_p(A)) = 3 + (33)^{\frac{1}{2}} = 8.745$ . Thus,  $\alpha(A) \leq \alpha(\gamma_p(A)) = 8.745$ . Actually, we have  $\alpha(A) = 8.657$ . Applying to  $A$  the inequality (v) of Sec. 1 with  $\phi = l_\infty, l_1$ , we obtain, respectively,

$$\alpha(A) \leq \gamma_{l_\infty}(A) = 16, \quad \alpha(A) \leq \gamma_{l_1}(A) = 10.$$

The upper bound  $\gamma_{l_2}(A)$  is more difficult to compute. It is the largest eigenvalue of  $\frac{1}{2}(A + A^*)$ . One obtains  $\alpha(A) \leq \gamma_{l_2}(A) = 9.039$ . It is interesting to note that we can obtain for  $\alpha(A)$  a better upper bound than those given by  $\gamma_{l_\infty}(A)$ ,  $\gamma_{l_1}(A)$ ,  $\gamma_{l_2}(A)$ , without even computing the spectral abscissa of  $\gamma_p(A)$ . Indeed, applying property (v) of Sec. 1 to  $\gamma_p(A)$  with  $\phi = l_1$ , we obtain

$$\alpha(A) \leq \alpha(\gamma_p(A)) \leq \gamma_{l_1}(\gamma_p(A)) = 9.$$

From Proposition 4.2 it follows that if, for a given  $A \in M_n(\mathbb{C})$ , the  $k \times k$  matrix  $\gamma_p(A)$  is stable for some equilibrated vectorial norm  $p$  of order  $k$  on  $\mathbb{C}^n$ , then  $A$  is also stable. Since  $\gamma_p(A)$  is essentially nonnegative, the following known theorem [8] will be very helpful.

**PROPOSITION A.** *Let  $S$  be an essentially nonnegative  $k \times k$  matrix, and let  $d_j$  denote the determinant of the  $j \times j$  principal submatrix of  $S$  lying in the first  $j$  rows and columns ( $j = 1, \dots, k$ ). Then the following statements are equivalent:*

- (1)  $-S$  is an  $M$ -matrix;
- (2)  $-d_1 > 0, d_2 > 0, -d_3 > 0, \dots, (-1)^k d_k > 0$ ;
- (3)  $S$  is stable.

(For other equivalent statements see [8] and [9]; see also [19] and [10, vol. 2, p. 74].)

**PROPOSITION 4.3.** *Let  $A \in M_n(\mathbb{C})$ , let  $p$  be any equilibrated vectorial norm of order  $k$  on  $\mathbb{C}^n$ , and let  $d_j$  be the determinant of the  $j \times j$  principal submatrix of  $\gamma_p(A)$  lying in the first  $j$  rows and columns ( $j = 1, \dots, k$ ). If*

$$-d_1 > 0, \quad d_2 > 0, \quad -d_3 > 0, \quad \dots, \quad (-1)^k d_k > 0,$$

*then  $A$  is stable.*

*Proof.* Since  $\gamma_p(A)$  is an essentially nonnegative  $k \times k$  matrix, the stability of  $\gamma_p(A)$  follows from Proposition A. Now, by Proposition 4.2,  $\alpha(A) < 0$ , i.e.,  $A$  is stable. ■

REMARK 4.3. By virtue of Proposition A, the conditions on the  $d_i$ 's in Proposition 4.3 can be replaced by the assumption that  $-\gamma_p(A)$  is an  $M$ -matrix.

EXAMPLE 4.2. Let

$$A = \begin{pmatrix} -12 & 1 & 6 & 2 \\ 1 & -8 & -4 & 4 \\ 6 & -4 & -14 & -3 \\ 2 & 4 & -3 & -14 \end{pmatrix},$$

and consider the partition  $\pi = \{0, 1, 2, 4\}$  of  $\mathbf{C}^4$ , which induces the indicated partitioning of  $A$ . Denoting  $p = (l_\infty, \pi)$ , we have

$$\gamma_p(A) = \begin{pmatrix} -12 & 1 & 8 \\ 1 & -8 & 8 \\ 6 & 4 & -11 \end{pmatrix},$$

and thus  $d_1 = -12 < 0$ ,  $d_2 = 95 > 0$ ,  $d_3 = -197 < 0$ . Consequently,  $A$  is stable. Note that with the partition  $\pi_0 = \{0, 4\}$  (i.e., no partition), we obtain only  $\alpha(A) \leq \gamma_{l_\infty}(A) = \gamma_{l_1}(A) = 1$ , while  $q = (l_\infty; \pi_1)$ , where  $\pi_1 = \{0, 2, 4\}$ , gives

$$\gamma_q(A) = \begin{pmatrix} -7 & 8 \\ 10 & -11 \end{pmatrix},$$

which is not stable. This latter matricial logarithmic derivative gives the upper bound  $\alpha(A) \leq \alpha(\gamma_q(A)) = -9 + (84)^{1/2} \approx 0.165$ .

For a matrix  $A = (a_{ij}) \in M_n(\mathbf{C})$ , we shall denote

$$\tilde{A} = \begin{pmatrix} \operatorname{Re} a_{11} & |a_{12}| & \cdots & |a_{1n}| \\ |a_{21}| & \operatorname{Re} a_{22} & \cdots & |a_{2n}| \\ \vdots & \vdots & \ddots & \vdots \\ |a_{n1}| & |a_{n2}| & \cdots & \operatorname{Re} a_{nn} \end{pmatrix}.$$

PROPOSITION 4.4 [6]. *Let  $A \in M_n(\mathbf{C})$ . Then  $\alpha(A) \leq \alpha(\tilde{A})$ .*

*Proof.* If in Proposition 4.2 we take  $p = (l_\infty, \pi)$ , where  $\pi$  is the partition determined by  $\{0, 1, 2, \dots, n\}$  (i.e., the finest partition of  $\mathbf{C}^n$ ), then we obtain  $\gamma_p(A) = \tilde{A}$ , and so  $\alpha(A) \leq \alpha(\tilde{A})$ . ■

PROPOSITION 4.5. Let  $A = (a_{ij}) \in M_n(\mathbb{C})$ , and denote

$$d_i = \begin{vmatrix} \operatorname{Re} a_{11} & |a_{12}| & \cdots & |a_{1i}| \\ |a_{21}| & \operatorname{Re} a_{22} & \cdots & |a_{2i}| \\ \vdots & \vdots & \ddots & \vdots \\ |a_{i1}| & |a_{i2}| & \cdots & \operatorname{Re} a_{ii} \end{vmatrix} \quad (i = 1, \dots, n).$$

If  $-d_1 > 0$ ,  $d_2 > 0$ ,  $-d_3 > 0, \dots$ ,  $(-1)^n d_n > 0$ , then  $A$  is stable.

*Proof.* This is an immediate consequence of Proposition 4.4 and Proposition A. ■

PROPOSITION 4.6 [16]. Let  $B$  be an essentially nonnegative matrix in  $M_n(\mathbb{R})$ . If  $A \in M_n(\mathbb{C})$  and  $\tilde{A} \leq B$ , then  $\alpha(A) \leq \alpha(B)$ .

*Proof.* We have, making use of Proposition 4.4,  $\alpha(A) \leq \alpha(\tilde{A}) \leq \alpha(B)$ . ■

PROPOSITION 4.8. Let  $A \in M_n(\mathbb{C})$ , let  $k \in \{1, 2, \dots, n\}$ , and let  $N_k$  denote the set of all equilibrated vectorial norms of order  $k$  on  $\mathbb{C}^n$ . Then

$$\alpha(A) = \inf_{p \in N_k} \alpha(\gamma_p(A)).$$

*Proof.* By Proposition 4.2,  $\alpha(A) \leq \alpha(\gamma_p(A))$  for all  $p \in N_k$ . Let  $\varepsilon > 0$ , and denote  $\delta = \varepsilon/3$ . There exists a nonsingular  $S \in M_n(\mathbb{C})$  such that  $T = SAS^{-1}$  is in the Jordan canonical form. Denoting  $D = \operatorname{diag}(1, \delta, \delta^2, \dots, \delta^{n-1})$ , we have

$$DTD^{-1} = \begin{pmatrix} \lambda_1 & 0 & 0 & \cdots & 0 & 0 \\ \delta_1 & \lambda_2 & 0 & \cdots & 0 & 0 \\ 0 & \delta_2 & \lambda_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_{n-1} & 0 \\ 0 & 0 & 0 & \cdots & \delta_{n-1} & \lambda_n \end{pmatrix},$$

where  $\delta_j \in \{0, \delta\}$  ( $j = 1, \dots, n-1$ ) and  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the eigenvalues of  $A$  (not necessarily distinct). Let  $\pi$  be the partition of  $\mathbb{C}^n$  defined by  $\{0, 1, 2, \dots, k-1, n\}$ , and let  $p = (l_\infty, \pi)$ . Clearly,  $p \in N_k$ . Now, it can be easily

seen that

$$\gamma_p(DTD^{-1}) = \begin{pmatrix} \operatorname{Re}\lambda_1 & 0 & 0 & \cdots & 0 & 0 \\ \delta_1 & \operatorname{Re}\lambda_2 & 0 & \cdots & 0 & 0 \\ 0 & \delta_2 & \operatorname{Re}\lambda_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \operatorname{Re}\lambda_{k-1} & 0 \\ 0 & 0 & 0 & \cdots & \delta_{k-1} & \beta \end{pmatrix},$$

where

$$\beta = \max_{j=k+1, \dots, n} \{ \operatorname{Re}\lambda_k, \operatorname{Re}\lambda_j + \delta_{j-1} \}.$$

Making use of property (v) of Sec. 1, we obtain

$$\alpha(\gamma_p(DTD^{-1})) \leq 2\delta + \max_{j=1, \dots, n} \operatorname{Re}\lambda_j < \alpha(A) + \varepsilon.$$

Denoting by  $p_0$  the DS-transform of the vectorial norm  $p$ , we have

$$\begin{aligned} \alpha(\gamma_{p_0}(A)) &= \alpha(\gamma_p(DSAS^{-1}D^{-1})) \\ &= \alpha(\gamma_p(DTD^{-1})) < \alpha(A) + \varepsilon. \end{aligned}$$

Consequently,  $\inf_{p \in N_k} \alpha(\gamma_p(A)) = \alpha(A)$ . ■

REMARK 4.4. For the special case  $k=1$ , Proposition 4.6 has been proved separately by Ström [22, 23] and Pao [18] (see also [7]).

## REFERENCES

- 1 W. A. Coppel, *Stability and Asymptotic Behavior of Differential Equations*, Heath, Boston, 1965.
- 2 G. Dahlquist, Stability and error bounds in the numerical integration of Ordinary Differential equations, in *Trans. Royal Inst. Technol.* **130**, Stockholm, 1959.
- 3 E. Deutsch, Matricial norms, *Numer. Math.* **16** (1970), 73–84.
- 4 E. Deutsch, On matricial norms subordinate to vectorial norms, *Math. Z.* **122** (1971), 142–150.

- 5 E. Deutsch, On vectorial norms and pseudonorms, *Proc. Amer. Math. Soc.* **28** (1971), 18–24.
- 6 E. Deutsch, The spectral abscissa of partitioned matrices, *J. Math. Anal. Appl.* **50** (1975), 66–73.
- 7 E. Deutsch, On matrix norms and logarithmic norms, *Numer. Math.* **24** (1975), 49–51.
- 8 K. Fan, Topological proofs for certain theorems on matrices with non-negative elements, *Monatsh. Math.* **62** (1958), 219–237.
- 9 M. Fiedler and V. Pták, On matrices with non-positive off-diagonal elements and positive principal minors, *Czech. Math. J.* **12** (87) (1962), 382–400.
- 10 F. R. Gantmacher, *The Theory of Matrices*, Chelsea, New York, 1959.
- 11 G. A. Hewer, A generalization of an inequality of Coppel, *Proc. Amer. Math. Soc.* **44** (1974), 151–156.
- 12 A. S. Householder, *The Theory of Matrices in Numerical Analysis*, Blaisdell, New York, 1964.
- 13 L. V. Kantorovič, The method of successive approximations for functional equations, *Acta Math.* **71** (1939), 63–97.
- 14 G. Köthe, *Topological Vector Spaces I*, Springer, New York, 1969.
- 15 S. M. Lozinskii, Error estimates for the numerical integration of ordinary differential equations—I, *Izv. Vyssh. Uchebn. Zaved. Mat.* No. 5(6) (1958), 52–90; errata No. 5(12) (1959), 222.
- 16 S. M. Lozinskii, A contribution to the theory of finite matrices, *Sov. Math.—Dokl.* **163** (1965), 1014–1016.
- 17 A. M. Ostrowski, On some metrical properties of operator matrices and matrices partitioned into blocks, *J. Math. Anal. Appl.* **2** (1961), 161–209.
- 18 C. V. Pao, Logarithmic derivatives of a square matrix, *Linear Algebra Appl.* **6** (1973), 159–164.
- 19 G. Poole and T. Boullion, A survey on  $M$ -matrices, *SIAM Rev.* **16** (1974), 419–427.
- 20 F. Robert, Normes vectorielles de vecteurs et de matrices, *Rev. Fr. Trait. Inf. Chiffres* **7** (1964), 261–299.
- 21 F. Robert, Sur les normes vectorielles régulières sur un espace vectoriel de dimension finie, *C. R. Acad. Sci. Paris* **261** (1965), 5173–5176.
- 22 T. Ström, On logarithmic norms, Report NA 69.06, Dept. of Inf. Process. Comput. Sci., The Royal Inst. of Technol., Stockholm, 1969.
- 23 T. Ström, On logarithmic norms, *SIAM J. Numer. Anal.* **12** (1975), 741–753.
- 24 R. S. Varga, *Matrix Iterative Analysis*, Prentice-Hall, Englewood Cliffs, 1962.

*Received 20 July 1976; revised 20 September 1976*